

Full Length Research Paper

Some Aspects on the Reproductive Biology of Greater Forkbeard Phycis Blennoides (Brünnich, 1768) in Western Algerian Coasts (Osteichthyes, Gadidae)

Sofiane Med El Amine Benghali^{1*}, Salim Mouffok², Ali Kherraz¹, Zitouni Boutiba²

¹ Department of Marine Science and Aquaculture, FSNV-UMAB-Mostaganem (Algeria)

² Environmental Surveillance Laboratory, Department of Biology, Faculty of Natural Sciences and Life, University Oran (Algeria)

Received May 30, 2013; Accepted July 20 2014

ABSTRACT

The aim of this work was to study the annual reproductive cycle of the greater forkbeard (*Phycis blennoides*, Brünnich, 1768) through analysis and description of some physiological index. A sample of 225 females was captured between January 2012 and December 2012 from the Western coast of Algeria, from Mostaganem. Fish length and weight varied between 17.60-37.80 cm and 38.33-403.03 g, respectively. Condition factors (K), gonadosomatic index (G.S.I.) and hepatosomatic index (H.S.I.) were calculated monthly.

Sex ratio of males to females was 1:0.91. The estimated length where 50% of analyzed individuals were sexual maturity was 24.73 cm for females. The peak value of the gonad somatic index was recorded in September and continued throughout October, indicating the highest spawning activity when the Kn values are low.

Key words: Greater Forkbeard, *Phycis Blennoides*, Gonadosomatic Index, Hepatosomatic Index, First Maturity, Spawning Period

1. INTRODUCTION

The greater forkbeard, *Phycis blennoides* (Brünnich, 1768) is a common gadoid occurring in the Mediterranean and in the North-eastern Atlantic, from Iceland to Morocco (Tortonese, 1975; Fisher *et al.*, 1987; Davis & Edwards, 1988; Whitehead *et al.*, 1989). It is usually found on muddy or sand bottoms in depths of 100-650m, the total length at first sexual maturity for males and females are 18-20 cm and 22-23 cm

respectively (Gallardo-Cabello & Gual-Frau, 1984). In the Mediterranean, *P. blennoides* spawns from June to August (Gallardo-Cabello & Gual-Frau, 1984).

Taking into account that length at maturity, fecundity and sex ratio are some of the most important parameters in studying reproductive dynamics of gadoids population, this study was carried out by examination of annual changes of the gonado-somatic index

*Corresponding author: sbenghali@yahoo.com

(GSI), hepato-somatic index (HSI) and condition factor (Kn) in order to evaluate the level of the exploitation in the Algerian coasts.

In fact, on southern Mediterranean coast, the knowledge on *P. blennoides* is still fragmentary, being limited to some remarks about Algerian waters.

Therefore, the aim of the present paper was to study the reproduction and condition of this species. This is the first paper with a complete analysis of the biology of *P. blennoides* in the south-western Mediterranean.

2. MATERIAL AND METHODS

Reproduction study

A series of biological samples was conducted on specimens of *P. Blennoides* caught by the commercial trawlers in the sampling area of Mostaganem (Fig.1). Specimens' total length was measured and both sex and maturity were reported for females which represent the focus of our study, four different stages were defined (Table 1) according to (ICES, 2007).

For each fish, total length (TL) was measured using a simple caliper to the nearest 0.1 cm and weighed (Wt) to the nearest 0.1 g. Fishes were gutted, and gonads were removed and weighed (Wg) with three decimal accuracy. Sex was determined by macroscopic observation of the gonads (Macer, 1974). Sex ratio was examined using χ^2 (Chi-square) test with a probability level of 0.05 to test differences in relation to the expected ratio 1:1. The gonadosomatic index (GSI) was estimated as: $GSI = Wg / Wt \times 100$. To estimate size at first sexual maturity, the data were fitted in equation: $P = 1/(1+e^{(abxL)})$; where P is probability that individuals are sexually matured and L is their length. The length when 50% of analyzed individuals were mature was

calculated according to Sparre and Venema (1998): $L50\% = a/b$.

Finally, to monitor morphological variations, the condition index was calculated to assess the degree of overweight consecutive to genital development and repletion state of the target species. Condition factor was studied in females in order to show differences of Kn (Le Cren, 1951) related to time, according to the formula: $Kn = W/Wth$ with $Wth = aL^b$ where "W" is the total weight, "Wth" is the theoretical weight, "a" and "b" are coefficients of the relative growth between weight and length and "L" is total length.

3. RESULT

Study of reproduction

1) Sex-ratio

After sexing of 461 specimens we found a sampling rate of 51.30% of Males significantly more important than females sex ratio (48.70%) the sex ratio (male: female) was 1:0.91 and it was statistically different from 1:1 ratio ($X^2=24.37$; $P<0.05$) (Table 2).

The length abundance curve is shown in Fig. 2. Fig. 3 shows a variation of the percentage of females per month. The female's percentage is dominant during fall and spring season, declining in the winter and summer period.

This finding indicates that males are, on average, significantly larger than females. As regards the sexual maturation of females, different stages of maturation of the gonads during different months of the year are shown in Fig. 4.

2) Indices of fish condition

In our study we have used three indexes to determinate the spawning period of the species in the study area: the gonado-somatic index (GSI), hepatosomatic index (HSI) and condition index (Kn). These allowed to

quantify morphological changes of the specimens and to identify reproduction period by studying the evolution of maturity stages of the ovary.

Hepato-Somatic Index (HSI) and Gonado-Somatic Index (GSI) and condition factor (Kn).

Monthly averages of GSI and HSI calculated from 225 females are plotted in Fig. 5. Only one peak were observed corresponding to the maximum annual spawning period of the population. The highest value of GSI were found in September (0.48 ± 0.06) and the lowest values in November (0.06 ± 0.04).

The highest values of the HSI occurred in July (9.19 ± 0.87) and the lowest fall in November (2.53 ± 0.34) (Fig. 5). Fig. 6 shows the condition factor Kn by seasons in both sexes. The values of Kn resulted overweight, thus revealing breeding events and confirming a rapid maturation occurring from July (1.12 ± 0.11) to September (0.91 ± 0.15) when the values of Kn are very low with irregular variations.

3) Length at maturity

For the statistical method, the L50 point estimated the body size at sexual maturity at 24.73 cm (Fig. 7). All data are combined in Table 3. Our results confirmed values reported for Mediterranean.

Fisheries which differ from those from the North Atlantic where specimen's length at maturity is longer than that they found in the Mediterranean Sea. Total individuals' length of the monthly samples ranged from a minimum of **17.50** centimeters to a maximum of **43.5** cm. Minimum sizes correspond to females and maximum sizes correspond to males.

4. DISCUSSION

This study presents first data of reproductive characteristics of forkbeard in the south-western Mediterranean Sea and therefore results were compared with other gadiform species common for this area.

We found a change in the rate of femininity with a significant dominance during the fall, which seems to correspond with the period when we recorded the maximum peak of the RGS which could correspond to a strong and early maturation of the ovaries.

The sex ratio showed a predominance of males, the catch rate (48.70%) for females (51.30%) for males, the sex ratio (male: female) is 1:0.91 In fact, this value is significantly different ($\chi^2 = 24.37$, $p > 0.05$) of the 1:1 theoretical value. The study of sex ratio depending on the size shows the dominance of male individuals up to size 26 cm, and that beyond this size the proportions of females are dominant, to the size where all females reach sexual maturity. These results are in agreement with the different regions of the Mediterranean (Gordon *et al.*, 1995. Rotllant *et al.*, 2002.). This predominance of females in older individuals could be explained by several authors by availability or larger female catchability; is a higher natural mortality in males. These studies showed that the females grow faster than males: in four years (Cohen *et al.*, 1990.).

The RGS is a real coefficient of gonadal maturation. Its increase coincides with gametogenesis while its decrease indicates an active spawning (Lahaye, 1972). Tracking monthly changes RGS allowed us to know the times of sexual activity *P. blennoides* and its breeding season.

We observed the presence of gravid females with a maximum of 0.48 RGS September also, there are resting

females in summer and winter; these observations could be explained by the fact that after spawning adult females would regain deep waters. Similarly, males have emissions of their sexual products with a maximum RGS 0.50 in September, this shows that the eggs are laid at the studied species could take place in autumn. Our results are almost similar to those obtained by other authors. They all define clearly the spawning period in autumn; This situation is common to all the Mediterranean coast (Gordon and Duncan, 1985) and in the Ionian Sea (Matarresse *et al.*, 1998). Presence of ripe females indicated that spawning of *Phycis blennoides* occurs during early autumn to early winter. Rotllant *et al.*, (2002) investigated population of *Phycis blennoides* in the western Mediterranean Sea. Mature females in their study were found only in autumn. Parallel to the RGS, we studied the RHS since the energy required for gonad maturity comes from fat reserves stored in the liver. The observation of the temporal evolution of the hepatosomatic report *Phycis blennoides* females showing phases of hepatic synthesis and consumption phases of liver lipids. The largest decline RHS is during the autumn which synchronizes perfectly with the period of mass reproduction in females and coincides with the transfer of liver reserves to the gonad.

The results of the study of the condition index (Kn) in females, set and confirm the spawning period in the range from September to November, as from September, the value of Kn begins to decrease with a minimum in October, which could be explained by a loss of organic matter associated with the laying period.

We estimated the size at first maturity (L50) to 24.73 cm. According to

Rodriguez - Cabello *et al.*, (1998), the size at first maturity of females of the Mediterranean is smaller than that found in the North Atlantic; this difference (also found in males) was explained by suggesting a possible relationship between the maturity of the species and latitude (Lam, 1983). Our findings and conclusions support obtained by various authors mentioned above suggests that the reproductive parameters of *Phycis blennoides* differ from one region to another, probably under the influence of various environmental and geographical parameters (Leloup & Oliverau, 1951 Relini & Orsi - Relini 1987 Capapé *et al.*, 1991; Demestre & Martin, 1993; Guijarro *et al.*, 2007), as the passage of Atlantic currents entering the Mediterranean through the Strait Gibraltar rich in organic matter providing an ideal enrichment Algerian deep waters (Cartes *et al.*, 2002), Silva (1986) calculated length at first maturity at 41 cm for females and 36 cm for males in Azorean waters and in this study that length was lower and equal to 24.73 cm for females.

In the conclusion, this study shows that Greater forkbeard has group-synchronous ovarian development and spawning season from early September to late November. The obtained results from this study are important input data for management and stock assessment of this commercially important fish species.

5. REFERENCES

- Alonso-Fernandez, A., Dominguez-Petit, R., Bao, M., Rivas, C. and Saborido-Ray, F. 2008. Spawning pattern and reproductive strategy of female pouting *Trisopterus luscus* (Gadidae) on the Galician shelf of north-western Spain. *Aquat. Living Resour.*, 21:383-393. doi:10.1051/alr:2008059
- Beverton R.J.H & Holt S.J., 1957. On the dynamics of exploited fish populations. *Fishery investigation*, series II, 19:1-533
- Bilgin, S., Bal, H. and Tasci, B. 2012. Length based growth estimation and reproduction biology of whiting, *Merlangius merlangus* (Nordman, 1840) in the Southeast Black Sea. *Turk. J. Fish. Aquat. Sc.*, 12:871-881. doi:10.4194/1303-2712-v12-4-15
- Capapé C., Tomasini J.A & Bouchereau J.L., 1991. Observations sur la biologie de reproduction de la petite roussette, *Scyliorhinus canicula* (Linnaeus, 1758) (Pisces, Scyliorhinidae du golfe du Lion (France méridionale). *Ichthyophysiological Acta*, 13:87-109
- Cartes J.E., Abelló P., Lloris D., Carbonell A., Torres P., Maynou F. & Gil De Sola L., 2002. Feeding guilds of western Mediterranean demersal fish and crustaceans: an analysis based on a spring survey. *Scientia Marina*, 66 (Suppl. 2):209-220
- Cohen D. M., Inada T., Iwamoto T., Scialabba N., 1990 – FAO Species catalogue. Gadiform fishes of the world. *Fisheries Synopsis* 125/10:442 pp.
- Edwards, P., R. S. V. Pullin, and J. A. Gartner. 1988. Research and development of integrated crop-livestock-fish farming systems in the tropics. *ICLARM Studies and Reviews* 16, 53 pp. International Center for Living Aquatic Resources Management, Manila, Philippines.
- Demestre, M. and P. Martin. 1993. Optimum exploitation of a demersal resource in the western Mediterranean: the fishery of the deep-water shrimp *Aristeus antennatus* (Risso, 1916). *Sci. Mar.*, 57:175-182
- Fischer, W., M. Schneider & M.L. Bauchot. 1987. Fiches FAO d'identification des espèces pour les besoins de la pêche: Méditerranée et Mer Noire (zone de pêche 37). II Vertébrés (FAO species identification sheets for fishery purposes: Mediterranean and Black Sea (fishing area 37). II Vertebrates). FAO, Rome, pp. 761-1530
- Gallardo-Cabello M., 1986b - Analisis de las frecuencias de talla por medio de los métodos de Petersen, Cassie y Bhattacharya, para la determinación de la edad de la brotola *Phycis blennoides* (Brünnich, 1768) en el Mediterráneo Occidental (Pisces: Gadidae). *Ann. Inst. Cienc. Mar Limnol. Univ. nat. autón. México*, 13:187-196
- Gordon, J.D.M. and Duncan, J.A.R. 1985b. The biology of fish of the Family Moridae in the deep-water of the Rockall Trough. *Journal of the Marine Biological Association of the United Kingdom* 65:475-485

- Gordon, J.D.M., Merrett, N.R. and Haedrich, R.L. 1995. Environmental and biological aspects of slope-dwelling fishes of the North Atlantic. pp. 1-26 in Hopper, A.G., editor. Deep-Water Fisheries of the North Atlantic Oceanic Slope. Kluwer Academic Publishers, Netherlands
- Guijarro B., Massutí E., Moranta J. & Díaz P., 2007. Population dynamics of the red-shrimp *Aristeus antennatus* in the Balearic Islands (western Mediterranean): short spatio-temporal differences and influence of environmental factors. *Journal of Marine Systems*, 71:385-402
- Gulland J.A., 1983. Fish stock assessment. A manual of basic methods. Ed. John Wiley and Sons, Chichester, UK, FAO/Wilies Series on Food and Agriculture, Vol.1., 223 pp.
- Gutiérrez-Estrada, J. C., Pulido-Calvo, I. and Prenda, J. 2000. Gonadosomatic index estimates of an introduced pumpkinseed (*Lepomis gibbosus*) population in a Mediterranean stream, using computational neural networks. *Aquat. Sci.*, 62:350-363. doi:10.1007/PL00001340
- Kirkwood G.P, Aukland R. & Zara J.S., 2001. Length Frequency Distribution Analysis (LFDA), Version 5.0. MRAG Ltd., London, UK.
- Lahaye (J.), 1972. Cycles sexuels de quelques poissons plats des côtes bretonnes. *Rev. Trav. Inst. Pêches marit.*, 36 (2):191-207
- Lam T., 1983. Environmental influences of gonadal activity in fish. In: Hoar W.S., Randall D.J. & Donaldson E.M. (eds.): *Fish Physiology*, 9. Academic Press, New York. London, Part B, pp. 65-116
- Le Cren E.D., 1951. The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *Journal of Animal Ecology*, 20: 201-219
- Leloup J & Olivereau M., 1951. Données biométrique comparatives sur la roussette (*Scyllium canicula* L.) de la Manche et de la Méditerranée. *Vie et Milieu*, 2:182-209
- Macer, C.T. 1974. The reproductive biology of the horse mackerel *Trachurus trachurus* (L.) in the North Sea and English Channel. *J. Fish Biol.*, 6:415-438. doi:10.1111/j.1095-8649.1974.tb04558.x
- Matarrese, A., D'Onghia, G., Basanisi, M. and Mastrototaro F. 1998. Spawning and recruitment of *Phycis blennoides* (Phycidae) from the north-western Ionian Sea (middle-eastern Mediterranean). *Ital. J. Zool.*, 65:203-209. doi:10.1080/11250009809386814
- Metin, G., Ilkyaz, A.T. and Kinacigil H.T. 2008. Growth, Mortality, and Reproduction of Poor Cood (*Trisopterus minutus* Linn., 1758) in the Central Aegean Sea. *Turk. J. Zool.*, 32:43-51. doi:10.1017/S0025315410000147
- Murua, H. and Motos, L. 2006. Reproductive strategy and spawning activity of the European hake *Merluccius merluccius* (L.) in the Bay of Biscay. *J. Fish Biol.*, 69:1288-1303. doi:10.1111/j.1095-8649.2006.01169.x

- Pauly D. & Moreau J., 1997. Méthodes pour l'évaluation des ressources halieutiques. Cépaduès-Editions, Toulouse, pp 288
- Relini G. & Orsi-Relini L., 1987. The decline of red shrimps stocks in the Gulf of Genova. *Investigación Pesquera*, 51:245-260
- Rodríguez-Cabello C., Velasco F. & Olaso I., 1998. Reproductive biology of lesser spotted dogfish *Scyliorhinus canicula* (L., 1758) in the Cantabrian Sea. *Scientia Marina*, 62:187-191
- Rotllant, G., Moranta, J., Massuti, E., Sarda, F. and Morales-Nin, B. 2002. Reproductive biology of three gadiforms fish species through the Mediterranean deep-sea range (147-1850 m). *Sci. Mar.*, 66:157-166
- Šantić, M., Pallaoro, A., Stagličić, N., Mikulandra, I. and Jardas I. 2010. Covariation of gonadosomatic index, condition factor and length-weight relationship of poor cod, *Trisopterus minutus* (Gadidae), in the Eastern Adriatic Sea. *Cybium*, 34:279-284
- Silva, H.M. 1986. Reproduction of the forkbeard *Phycis phycis* (Linnaeus, 1766) in Azorean waters. *ICES CM* 1986/G:60
- Sparre P. & Venema S.C., 1996. Introduction à l'évaluation des stocks des poissons tropicaux. Première partie: Manuel FAO. Document Technique sur les Pêches, 306 (1), Danida, Romepp, 401
- Statsoft INC. (2001). STATISTICA (data analysis software system), version 6, www.statsoft.com
- Sparre, P. and Venema, S.C. 1998. Introduction to tropic fish stock assessment. Part 1. Manual. FAO Fisheries Technical Paper.No. 306/1, Rev.2. FAO, Rome, pp. 407
- Tortonese, E. 1975. Osteichthyes (Pesci ossei), Parte Seconda. In: Fauna d'Italia XI., Edizioni Calderini, Bologna 636 pp
- Tsikliras, A.C., E. Antonopoulou & K.I. Stergiou. 2010. Spawning period of Mediterranean marine fishes. *Rev. Fish Biol. Fish.*, 20:499-538
- Ursin E., 1967. A mathematical model of some aspects of fish growth, respiration and mortality. *Journal of Fisheries Research Board of Canada*, 24:2355-2453.
- Von Bertalanffy L., 1938. A quantitative theory of organic growth. *Human Biology*, 10:181-213
- Von Bertalanffy, L. 1938. A quantitative theory of organic growth. *Hum. Biol.* 10:181-213
- Whitehead PJP. 1985. FAO species catalogue. Vol. 7. Clupeoid fishes of the world. Part 1. Chirocentridae, Clupeidae and Pristigasteridae. Rome, Italy. FAO Fish. Synop. 125, Vol. 7, Pt. 1, pp. 303

6. FIGURES



Figure 1: Study Area




Figure 2: Abundance curve. Results of χ^2 test show a predominance of one sex over the other by length of specimens (* $p < 0.05$)




Figure 3: Distribution of males and females of *Phycis blennoides* by season. χ^2 test results show a prevalence of one sex over the other per sampling month




Figure 4: Percentages of different stages of sexual maturity in *Phycis blennoides* females per month




Figure 5: Monthly trend of GSI and HSI with standard errors in *Phycis blennoides*




Figure 6: Condition index (Kn) with standard error according to the season in *Phycis blennoides* females




Figure 7: Size of first sexual maturity in *Phycis blennoides*

7. TABLES

MATURITY STAGES	DESCRIPTION
1. Immature/Resting	Small ovaries, with firm consistence and minimal visualization, transparent or pink grey, without opaque or hyaline oocytes.
2. Developing/Maturing	Medium or large ovaries, pink or yellow to orange, with visualization variable, present and obvious. Opaque oocytes present but without hyaline oocytes.
3. Spawning	Hydrated-Large ovaries, with firm consistence and visualization, pink or reddish orange. Opaque and hyaline oocytes present.
4. Post-spawning	Small or medium ovaries, flaccid, dark pink, orange or purple. Opaque and hyaline oocytes absent or residual.

Table 1: Different stages of maturity of *P. Blennoides* females

Sex	Total	Percentage
Females	225	48.70%
Males	236	*51.30%
Total	461	100%

Table 2: Percentage of sexes in *P. Blennoides* (*p<0,05).

Authors	Area	Males (cm)	Females (cm)
Cohen <i>et al.</i> , 1990	Atlantic	18	33
Rotllant <i>et al.</i> , 2002	Mediterranean	19.32	20
Present work 2014	West Algeria	*	24.73

Table 3: Summary of first sexual maturity length (L50) of *P. blennoides* females and males from different areas (* only females were studied to determine the size at sexual maturity)